



Monica Hum MD • FACS • FASCRS
Tonya Jordan MD
Joseph Mareno Jr. MD

Moviprep Preparation Instructions

Five days before the procedure

- If you are taking **Aspirin** based products or **Vitamin E**, please contact our office for further instructions. If you are currently taking blood thinners such as Coumadin or Plavix, please let us know immediately, as you must stop this medication several days prior to the exam to prevent excessive bleeding. You must also clear this with your primary care physician.

The entire day before the procedure

You may follow these directions or the manufacturer's directions as listed on their website:

http://www.salix.com/products/products_moviprep.asp

- You may have a light, low fiber breakfast early in the morning (eg. eggs, white toast – no butter). No vegetables, fruits, nuts, butter, milk, cheese, meat, or whole grain cereals.
- YOU MUST DRINK PLENTY OF CLEAR LIQUIDS THROUGHOUT THE ENTIRE DAY – it is recommended that you drink a minimum of 32 fluid ounces of clear liquids spread out over the day BEFORE you start your prep.
- You will be on a clear liquid diet. A clear liquid is any liquid that you can put in a glass and see straight through it. You may have Jello, bouillon, consommé, apple juice, grape juice, cranberry juice, Gatorade, black coffee, tea, cola, or water. Anything that melts into a clear liquid (sorbet, Italian ice, ice, lifesavers) is allowed. Milk, cream, anything with pulp, or alcoholic beverages is **NOT** allowed. Solid food is **NOT** allowed.

The evening before the procedure

- Mix the solution as directed; add one Pouch A & B into the bottle provided and add lukewarm drinking water to top of line on the bottle. Cap the bottle and shake to dissolve the powder.
- Beginning anytime between 4 & 7 p.m., start drinking 8 ounces of the fluid every fifteen to twenty minutes. Chilling the liquid or sipping through a straw makes it easier. You will finish the entire liter of liquid.
- Two hours after you finished the first liter, repeat the above steps with the other Pouch A & B, creating and drinking a second liter.
- Afterwards, you may then drink all the clear liquids you like. You are to STOP EATING OR DRINKING eight hours prior to your procedure time.

The morning of the procedure

- Patients taking daily medications may take their regular dosages the morning of the exam with very small sips of water.
- **For COLONOSCOPY:** report to the Endoscopy Lab located on the first floor of the 77 Bldg. Park in the NORTH parking deck. You should arrive at least one hour prior to your scheduled appointment time. The entire length of time you will spend at the hospital for colonoscopy will be approximately 2-3 hours.
- **For AMBULATORY SURGERY:** report to the McDonnell Surgical Center on Collier Road. Park in the McDonnell Surgical Center parking lot. You should arrive two hours prior to your scheduled surgery time.
- **For INPATIENT SURGERY:** report to the Admissions Desk in the main hospital (this is near elevator A). You should arrive two hours prior to your scheduled surgery time.



Monica Hum MD • FACS • FASCRS
Tonya Jordan MD
Joseph Mareno Jr. MD

Moviprep Preparation Instructions (cont.)

After the procedure

- You must have someone responsible drive you home after colonoscopy & ambulatory surgery - these are hospital rules - because you will be given a sedative for the procedure. Your procedure will be cancelled if you do not have a responsible escort home.
- Biopsy/pathology reports take about 3-4 business days to get from the laboratory. Please allow this much time before contacting our office for your results.
- If you have any questions, please contact our office prior to your exam.

Be sure to visit our website www.atlcrs.com to check our Frequently Asked Questions (FAQs) section.